

Legal Disclaimer: Clinical practice guidelines made available by WellCare are informational in nature and are not a substitute for the professional medical judgment of treating physicians or other health care practitioners. These guidelines are based on information available at the time and may not be updated with the most current information available at subsequent times. Individuals should consult with their physician(s) regarding the appropriateness of care or treatment options to meet their specific needs or medical condition. Disclosure of clinical practice guidelines is not a guarantee of coverage. Members of WellCare health plans should consult their individual coverage documents for information regarding covered benefits. WellCare does not offer medical advice or provide medical care, and therefore cannot guarantee any results or outcomes. WellCare does not warrant or guarantee, and shall not be liable for any deficiencies in the information contained herein or for any inaccuracies or recommendations made by independent third parties from whom any of the information contained herein was obtained.

Each child and family is unique; therefore, these Recommendations for Preventive Pediatric Health Care are designed for the care of children who are receiving competent parenting, have no manifestations of any important health problems, and are growing and developing in satisfactory fashion. Additional visits may become necessary if circumstances suggest variations from normal.

Developmental, psychosocial, and chronic disease issues for children and adolescents may require frequent counseling and treatment visits separate from preventive care visits.

These guidelines represent a consensus by the American Academy of Pediatrics (AAP) and Bright Futures. The AAP continues to emphasize the great importance of **continuity of care** in comprehensive health supervision and the need to avoid **fragmentation of care**.

The recommendations in this statement do not indicate an exclusive course of treatment or standard of medical care. Variations, taking into account individual circumstances, may be appropriate.

Copyright © 2008 by the American Academy of Pediatrics.

No part of this statement may be reproduced in any form or by any means without prior written permission from the American Academy of Pediatrics except for one copy for personal use.

Age ^a	INFANCY									EARLY CHILDHOOD						MIDDLE CHILDHOOD						ADOLESCENCE												
	Prenatal ^b	Newborn ^c	3-5 dy ^d	1 mo	2 mo	4 mo	6 mo	9 mo	12 mo	15 mo	18 mo	24 mo	30 mo	3 y	4 y	5 y	6 y	7 y	8 y	9 y	10 y	11 y	12 y	13 y	14 y	15 y	16 y	17 y	18 y	19 y	20 y	21 y		
HISTORY																																		
Initial/interval	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	
MEASUREMENTS																																		
Length/height and weight	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	
Head circumference	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	
Weight for length	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	
Body mass index																																		
Blood pressure ^e	★	★	★	★	★	★	★	★	★	★	★	★	★	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●		
SENSORY SCREENING																																		
Vision		★	★	★	★	★	★	★	★	★	★	★	★	●	●	●	●	★	●	★	●	★	●	★	●	★	●	★	●	★	●	★	●	
Hearing		● ^g	★	★	★	★	★	★	★	★	★	★	★	★	●	●	●	★	●	★	●	★	●	★	●	★	●	★	●	★	●	★	●	
DEVELOPMENTAL/BEHAVIORAL ASSESSMENT																																		
Developmental screening ^h													●	●																				
Autism screening ⁱ																																		
Developmental surveillance ^j																																		
Psychosocial/behavioral assessment																																		
Alcohol and drug use assessment																																		
PHYSICAL EXAMINATION^k																																		
	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●
PROCEDURES^l																																		
Newborn metabolic/hemoglobin screening	←	●	→																															
Immunization ^m		●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●
Hematocrit or hemoglobin						★			●		★	★		★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★
Lead screening ⁿ																																		
Tuberculin test ^o				★			★		★		★	★		★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★
Dyslipidemia screening ^p																																		
STI screening ^q																																		
Cervical dysplasia screening ^r																																		
ORAL HEALTH^s								★	★	● ^t	● ^t	● ^t	● ^t	● ^t	● ^t																			
ANTICIPATORY GUIDANCE^u	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●

a If a child comes under care for the first time at any point on the schedule, or if any items are not accomplished at the suggested age, the schedule should be brought up to date at the earliest possible time.
b A prenatal visit is recommended for parents who are at high risk, for first-time parents, and for those who request a conference. The prenatal visit should include anticipatory guidance, pertinent medical history, and a discussion of benefits of breastfeeding and planned method of feeding per AAP statement "The Prenatal Visit" (2001) [URL: <http://aapolicy.aappublications.org/cgi/content/full/pediatrics;107/6/1456>].
c Every infant should have a newborn evaluation after birth, breastfeeding encouraged, and instruction and support offered.
d Every infant should have an evaluation within 3 to 5 days of birth and within 48 to 72 hours after discharge from the hospital, to include evaluation for feeding and jaundice. Breastfeeding infants should receive formal breastfeeding evaluation, encouragement, and instruction as recommended in AAP statement "Breastfeeding and the Use of Human Milk" (2005) [URL: <http://aapolicy.aappublications.org/cgi/content/full/pediatrics;115/2/496>]. For newborns discharged in less than 48 hours after delivery, the infant must be examined within 48 hours of discharge per AAP statement "Hospital Stay for Healthy Term Newborns" (2004) [URL: <http://aapolicy.aappublications.org/cgi/content/full/pediatrics;113/5/1434>].
e Blood pressure measurement in infants and children with specific risk conditions should be performed at visits before age 3 years.
f If the patient is uncooperative, rescreen within 6 months per AAP statement "Eye Examination and Vision Screening in Infants, Children, and Young Adults" (1996) [URL: <http://aapolicy.aappublications.org/cgi/registr/pediatrics;98/1/153.pdf>].
g All newborns should be screened per AAP statement "Year 2000 Position Statement: Principles and Guidelines for Early Hearing Detection and Intervention Programs" (2000) [URL: <http://aapolicy.aappublications.org/cgi/content/full/pediatrics;106/4/796>].
h Joint Committee on Infant Hearing. "Year 2007 position statement: principles and guidelines for early hearing detection and intervention programs." *Pediatrics*. 2007;120:898-921.
i AAP Council on Children With Disabilities, AAP Section on Developmental Behavioral Pediatrics, AAP Bright Futures Steering Committee, AAP Medical Home Initiatives for Children With Special Needs Project Advisory Committee. Identifying infants and young children with developmental disorders in the medical home: an algorithm for developmental surveillance and screening. *Pediatrics*. 2006;118:405-420 [URL: <http://aapolicy.aappublications.org/cgi/content/full/pediatrics;118/1/405>].
j Gupta VB, Hyman SL, Johnson CP, et al. Identifying children with autism early? *Pediatrics*. 2007;119:152-153 [URL: <http://pediatrics.aappublications.org/cgi/content/full/pediatrics;119/1/152>].
k At each visit, age-appropriate physical examination is essential, with infant totally unclothed, older child undressed and suitably draped.
l These may be modified, depending on entry point into schedule and individual need.
m Newborn metabolic and hemoglobinopathy screening should be done according to state law. Results should be reviewed at visits and appropriate retesting or referral done as needed.
n Schedules per the Committee on Infectious Diseases, published annually in the January issue of *Pediatrics*. Every visit should be an opportunity to update and complete a child's immunizations.
o See AAP *Pediatric Nutrition Handbook*, 5th Edition (2003) for a discussion of universal and selective screening options. See also Recommendations to prevent and control iron deficiency in the United States. *MMWR Recomm Rep*. 1996;47(RR-3):1-36.
p For children at risk of lead exposure, consult the AAP statement "Lead Exposure in Children: Prevention, Detection, and Management" (2005) [URL: <http://aapolicy.aappublications.org/cgi/content/full/pediatrics;116/4/1036>]. Additionally, screening should be done in accordance with state law where applicable.
q Perform risk assessments or screens as appropriate, based on universal screening requirements for patients with Medicaid or high prevalence areas.
r Tuberculosis testing per recommendations of the Committee on Infectious Diseases, published in the current edition of *Red Book: Report of the Committee on Infectious Diseases*. Testing should be done on recognition of high-risk factors.
s "Third Report of the National Cholesterol Education Program (NCEP) Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III) Final Report" (2002) [URL: <http://circ.ahajournals.org/cgi/content/full/106/25/3143>] and "The Expert Committee Recommendations on the Assessment, Prevention, and Treatment of Child and Adolescent Overweight and Obesity: Supplement to Pediatrics." Supplement to *Pediatrics*. In press.
t All sexually active patients should be screened for sexually transmitted infections (STIs).
u All sexually active girls should have screening for cervical dysplasia as part of a pelvic examination beginning within 3 years of onset of sexual activity or age 21 (whichever comes first).
v Referral to dental home, if available. Otherwise, administer oral health risk assessment. If the primary water source is deficient in fluoride, consider oral fluoride supplementation.
w At the visits for 3 years and 6 years of age, it should be determined whether the patient has a dental home. If the patient does not have a dental home, a referral should be made to one. If the primary water source is deficient in fluoride, consider oral fluoride supplementation.
x Refer to the specific guidance by age as listed in Bright Futures Guidelines. (Hagan JF, Shaw JS, Duncan PM, eds. *Bright Futures: Guidelines for Health Supervision of Infants, Children, and Adolescents*. 3rd ed. Elk Grove Village, IL: American Academy of Pediatrics; 2008.)

● = to be performed ★ = risk assessment to be performed, with appropriate action to follow, if positive ← ● → = range during which a service may be provided, with the symbol indicating the preferred age

Recommended Immunization Schedule for Persons Aged 0 Through 6 Years—United States • 2009

For those who fall behind or start late, see the catch-up schedule

Vaccine ▼	Age ►	Birth	1 month	2 months	4 months	6 months	12 months	15 months	18 months	19–23 months	2–3 years	4–6 years
Hepatitis B ¹	HepB	HepB	HepB		<i>see footnote 1</i>	HepB						
Rotavirus ²				RV	RV	RV ²						
Diphtheria, Tetanus, Pertussis ³				DTaP	DTaP	DTaP	<i>see footnote 3</i>	DTaP				DTaP
<i>Haemophilus influenzae</i> type b ⁴				Hib	Hib	Hib ⁴		Hib				
Pneumococcal ⁵				PCV	PCV	PCV		PCV			PPSV	
Inactivated Poliovirus				IPV	IPV			IPV				IPV
Influenza ⁶								Influenza (Yearly)				
Measles, Mumps, Rubella ⁷								MMR		<i>see footnote 7</i>		MMR
Varicella ⁸								Varicella		<i>see footnote 8</i>		Varicella
Hepatitis A ⁹								HepA (2 doses)			HepA Series	
Meningococcal ¹⁰											MCV	

Range of recommended ages

Certain high-risk groups

This schedule indicates the recommended ages for routine administration of currently licensed vaccines, as of December 1, 2008, for children aged 0 through 6 years. Any dose not administered at the recommended age should be administered at a subsequent visit, when indicated and feasible. Licensed combination vaccines may be used whenever any component of the combination is indicated and other components are not contraindicated and if approved by the Food and Drug Administration for that dose of

the series. Providers should consult the relevant Advisory Committee on Immunization Practices statement for detailed recommendations, including high-risk conditions: <http://www.cdc.gov/vaccines/pubs/acip-list.htm>. Clinically significant adverse events that follow immunization should be reported to the Vaccine Adverse Event Reporting System (VAERS). Guidance about how to obtain and complete a VAERS form is available at <http://www.vaers.hhs.gov> or by telephone, 800-822-7967.

1. Hepatitis B vaccine (HepB). (Minimum age: birth)

At birth:

- Administer monovalent HepB to all newborns before hospital discharge.
- If mother is hepatitis B surface antigen (HBsAg)-positive, administer HepB and 0.5 mL of hepatitis B immune globulin (HBIG) within 12 hours of birth.
- If mother's HBsAg status is unknown, administer HepB within 12 hours of birth. Determine mother's HBsAg status as soon as possible and, if HBsAg-positive, administer HBIG (no later than age 1 week).

After the birth dose:

- The HepB series should be completed with either monovalent HepB or a combination vaccine containing HepB. The second dose should be administered at age 1 or 2 months. The final dose should be administered no earlier than age 24 weeks.
- Infants born to HBsAg-positive mothers should be tested for HBsAg and antibody to HBsAg (anti-HBs) after completion of at least 3 doses of the HepB series, at age 9 through 18 months (generally at the next well-child visit).

4-month dose:

- Administration of 4 doses of HepB to infants is permissible when combination vaccines containing HepB are administered after the birth dose.

2. Rotavirus vaccine (RV). (Minimum age: 6 weeks)

- Administer the first dose at age 6 through 14 weeks (maximum age: 14 weeks 6 days). Vaccination should not be initiated for infants aged 15 weeks or older (i.e., 15 weeks 0 days or older).
- Administer the final dose in the series by age 8 months 0 days.
- If Rotarix[®] is administered at ages 2 and 4 months, a dose at 6 months is not indicated.

3. Diphtheria and tetanus toxoids and acellular pertussis vaccine (DTaP). (Minimum age: 6 weeks)

- The fourth dose may be administered as early as age 12 months, provided at least 6 months have elapsed since the third dose.
- Administer the final dose in the series at age 4 through 6 years.

4. Haemophilus influenzae type b conjugate vaccine (Hib). (Minimum age: 6 weeks)

- If PRP-OMP (PedvaxHIB[®] or Comvax[®] [HepB-Hib]) is administered at ages 2 and 4 months, a dose at age 6 months is not indicated.
- TriHiBit[®] (DTaP/Hib) should not be used for doses at ages 2, 4, or 6 months but can be used as the final dose in children aged 12 months or older.

5. Pneumococcal vaccine. (Minimum age: 6 weeks for pneumococcal conjugate vaccine [PCV]; 2 years for pneumococcal polysaccharide vaccine [PPSV])

- PCV is recommended for all children aged younger than 5 years. Administer 1 dose of PCV to all healthy children aged 24 through 59 months who are not completely vaccinated for their age.

- Administer PPSV to children aged 2 years or older with certain underlying medical conditions (see *MMWR* 2000;49[No. RR-9]), including a cochlear implant.

6. Influenza vaccine. (Minimum age: 6 months for trivalent inactivated influenza vaccine [TIV]; 2 years for live, attenuated influenza vaccine [LAIV])

- Administer annually to children aged 6 months through 18 years.
- For healthy nonpregnant persons (i.e., those who do not have underlying medical conditions that predispose them to influenza complications) aged 2 through 49 years, either LAIV or TIV may be used.
- Children receiving TIV should receive 0.25 mL if aged 6 through 35 months or 0.5 mL if aged 3 years or older.
- Administer 2 doses (separated by at least 4 weeks) to children aged younger than 9 years who are receiving influenza vaccine for the first time or who were vaccinated for the first time during the previous influenza season but only received 1 dose.

7. Measles, mumps, and rubella vaccine (MMR). (Minimum age: 12 months)

- Administer the second dose at age 4 through 6 years. However, the second dose may be administered before age 4, provided at least 28 days have elapsed since the first dose.

8. Varicella vaccine. (Minimum age: 12 months)

- Administer the second dose at age 4 through 6 years. However, the second dose may be administered before age 4, provided at least 3 months have elapsed since the first dose.
- For children aged 12 months through 12 years the minimum interval between doses is 3 months. However, if the second dose was administered at least 28 days after the first dose, it can be accepted as valid.

9. Hepatitis A vaccine (HepA). (Minimum age: 12 months)

- Administer to all children aged 1 year (i.e., aged 12 through 23 months). Administer 2 doses at least 6 months apart.
- Children not fully vaccinated by age 2 years can be vaccinated at subsequent visits.
- HepA also is recommended for children older than 1 year who live in areas where vaccination programs target older children or who are at increased risk of infection. See *MMWR* 2006;55[No. RR-7].

10. Meningococcal vaccine. (Minimum age: 2 years for meningococcal conjugate vaccine [MCV] and for meningococcal polysaccharide vaccine [MPSV])

- Administer MCV to children aged 2 through 10 years with terminal complement component deficiency, anatomic or functional asplenia, and certain other high-risk groups. See *MMWR* 2005;54[No. RR-7].
- Persons who received MPSV 3 or more years previously and who remain at increased risk for meningococcal disease should be revaccinated with MCV.

Recommended Immunization Schedule for Persons Aged 7 Through 18 Years—United States • 2009

For those who fall behind or start late, see the schedule below and the catch-up schedule

Vaccine ▼	Age ►	7–10 years	11–12 years	13–18 years
Tetanus, Diphtheria, Pertussis ¹		see footnote 1	Tdap	Tdap
Human Papillomavirus ²		see footnote 2	HPV (3 doses)	HPV Series
Meningococcal ³		MCV	MCV	MCV
Influenza ⁴		Influenza (Yearly)		
Pneumococcal ⁵		PPSV		
Hepatitis A ⁶		HepA Series		
Hepatitis B ⁷		HepB Series		
Inactivated Poliovirus ⁸		IPV Series		
Measles, Mumps, Rubella ⁹		MMR Series		
Varicella ¹⁰		Varicella Series		

Range of recommended ages

Catch-up immunization

Certain high-risk groups

This schedule indicates the recommended ages for routine administration of currently licensed vaccines, as of December 1, 2008, for children aged 7 through 18 years. Any dose not administered at the recommended age should be administered at a subsequent visit, when indicated and feasible. Licensed combination vaccines may be used whenever any component of the combination is indicated and other components are not contraindicated and if approved by the Food and Drug Administration for that dose of

the series. Providers should consult the relevant Advisory Committee on Immunization Practices statement for detailed recommendations, including high-risk conditions: <http://www.cdc.gov/vaccines/pubs/acip-list.htm>. Clinically significant adverse events that follow immunization should be reported to the Vaccine Adverse Event Reporting System (VAERS). Guidance about how to obtain and complete a VAERS form is available at <http://www.vaers.hhs.gov> or by telephone, 800-822-7967.

1. Tetanus and diphtheria toxoids and acellular pertussis vaccine (Tdap). (Minimum age: 10 years for BOOSTRIX® and 11 years for ADACEL®)

- Administer at age 11 or 12 years for those who have completed the recommended childhood DTP/DTaP vaccination series and have not received a tetanus and diphtheria toxoid (Td) booster dose.
- Persons aged 13 through 18 years who have not received Tdap should receive a dose.
- A 5-year interval from the last Td dose is encouraged when Tdap is used as a booster dose; however, a shorter interval may be used if pertussis immunity is needed.

2. Human papillomavirus vaccine (HPV). (Minimum age: 9 years)

- Administer the first dose to females at age 11 or 12 years.
- Administer the second dose 2 months after the first dose and the third dose 6 months after the first dose (at least 24 weeks after the first dose).
- Administer the series to females at age 13 through 18 years if not previously vaccinated.

3. Meningococcal conjugate vaccine (MCV).

- Administer at age 11 or 12 years, or at age 13 through 18 years if not previously vaccinated.
- Administer to previously unvaccinated college freshmen living in a dormitory.
- MCV is recommended for children aged 2 through 10 years with terminal complement component deficiency, anatomic or functional asplenia, and certain other groups at high risk. See *MMWR* 2005;54(No. RR-7).
- Persons who received MPSV 5 or more years previously and remain at increased risk for meningococcal disease should be revaccinated with MCV.

4. Influenza vaccine.

- Administer annually to children aged 6 months through 18 years.
- For healthy nonpregnant persons (i.e., those who do not have underlying medical conditions that predispose them to influenza complications) aged 2 through 49 years, either LAIV or TIV may be used.
- Administer 2 doses (separated by at least 4 weeks) to children aged younger than 9 years who are receiving influenza vaccine for the first time or who were vaccinated for the first time during the previous influenza season but only received 1 dose.

5. Pneumococcal polysaccharide vaccine (PPSV).

- Administer to children with certain underlying medical conditions (see *MMWR* 1997;46[No. RR-8]), including a cochlear implant. A single revaccination should be administered to children with functional or anatomic asplenia or other immunocompromising condition after 5 years.

6. Hepatitis A vaccine (HepA).

- Administer 2 doses at least 6 months apart.
- HepA is recommended for children older than 1 year who live in areas where vaccination programs target older children or who are at increased risk of infection. See *MMWR* 2006;55(No. RR-7).

7. Hepatitis B vaccine (HepB).

- Administer the 3-dose series to those not previously vaccinated.
- A 2-dose series (separated by at least 4 months) of adult formulation Recombivax HB® is licensed for children aged 11 through 15 years.

8. Inactivated poliovirus vaccine (IPV).

- For children who received an all-IPV or all-oral poliovirus (OPV) series, a fourth dose is not necessary if the third dose was administered at age 4 years or older.
- If both OPV and IPV were administered as part of a series, a total of 4 doses should be administered, regardless of the child's current age.

9. Measles, mumps, and rubella vaccine (MMR).

- If not previously vaccinated, administer 2 doses or the second dose for those who have received only 1 dose, with at least 28 days between doses.

10. Varicella vaccine.

- For persons aged 7 through 18 years without evidence of immunity (see *MMWR* 2007;56[No. RR-4]), administer 2 doses if not previously vaccinated or the second dose if they have received only 1 dose.
- For persons aged 7 through 12 years, the minimum interval between doses is 3 months. However, if the second dose was administered at least 28 days after the first dose, it can be accepted as valid.
- For persons aged 13 years and older, the minimum interval between doses is 28 days.

The Recommended Immunization Schedules for Persons Aged 0 Through 18 Years are approved by the Advisory Committee on Immunization Practices (www.cdc.gov/vaccines/recs/acip), the American Academy of Pediatrics (<http://www.aap.org>), and the American Academy of Family Physicians (<http://www.aafp.org>).

DEPARTMENT OF HEALTH AND HUMAN SERVICES • CENTERS FOR DISEASE CONTROL AND PREVENTION

Catch-up Immunization Schedule for Persons Aged 4 Months Through 18 Years Who Start Late or Who Are More Than 1 Month Behind—United States • 2009

The table below provides catch-up schedules and minimum intervals between doses for children whose vaccinations have been delayed. A vaccine series does not need to be restarted, regardless of the time that has elapsed between doses. Use the section appropriate for the child's age.

CATCH-UP SCHEDULE FOR PERSONS AGED 4 MONTHS THROUGH 6 YEARS					
Vaccine	Minimum Age for Dose 1	Minimum Interval Between Doses			
		Dose 1 to Dose 2	Dose 2 to Dose 3	Dose 3 to Dose 4	Dose 4 to Dose 5
Hepatitis B ¹	Birth	4 weeks	8 weeks (and at least 16 weeks after first dose)		
Rotavirus ²	6 wks	4 weeks	4 weeks ²		
Diphtheria, Tetanus, Pertussis ³	6 wks	4 weeks	4 weeks	6 months	6 months ³
<i>Haemophilus influenzae</i> type b ⁴	6 wks	4 weeks if first dose administered at younger than age 12 months 8 weeks (as final dose) if first dose administered at age 12-14 months No further doses needed if first dose administered at age 15 months or older	4 weeks ⁴ if current age is younger than 12 months 8 weeks (as final dose) ⁴ if current age is 12 months or older and second dose administered at younger than age 15 months No further doses needed if previous dose administered at age 15 months or older	8 weeks (as final dose) This dose only necessary for children aged 12 months through 59 months who received 3 doses before age 12 months	
Pneumococcal ⁵	6 wks	4 weeks if first dose administered at younger than age 12 months 8 weeks (as final dose for healthy children) if first dose administered at age 12 months or older or current age 24 through 59 months No further doses needed for healthy children if first dose administered at age 24 months or older	4 weeks if current age is younger than 12 months 8 weeks (as final dose for healthy children) if current age is 12 months or older No further doses needed for healthy children if previous dose administered at age 24 months or older	8 weeks (as final dose) This dose only necessary for children aged 12 months through 59 months who received 3 doses before age 12 months or for high-risk children who received 3 doses at any age	
Inactivated Poliovirus ⁶	6 wks	4 weeks	4 weeks	4 weeks ⁶	
Measles, Mumps, Rubella ⁷	12 mos	4 weeks			
Varicella ⁸	12 mos	3 months			
Hepatitis A ⁹	12 mos	6 months			
CATCH-UP SCHEDULE FOR PERSONS AGED 7 THROUGH 18 YEARS					
Tetanus, Diphtheria/ Tetanus, Diphtheria, Pertussis ¹⁰	7 yrs ¹⁰	4 weeks	4 weeks if first dose administered at younger than age 12 months 6 months if first dose administered at age 12 months or older	6 months if first dose administered at younger than age 12 months	
Human Papillomavirus ¹¹	9 yrs	Routine dosing intervals are recommended ¹¹			
Hepatitis A ⁹	12 mos	6 months			
Hepatitis B ¹	Birth	4 weeks	8 weeks (and at least 16 weeks after first dose)		
Inactivated Poliovirus ⁶	6 wks	4 weeks	4 weeks	4 weeks ⁶	
Measles, Mumps, Rubella ⁷	12 mos	4 weeks			
Varicella ⁸	12 mos	3 months if the person is younger than age 13 years 4 weeks if the person is aged 13 years or older			

1. Hepatitis B vaccine (HepB).

- Administer the 3-dose series to those not previously vaccinated.
- A 2-dose series (separated by at least 4 months) of adult formulation Recombivax HB[®] is licensed for children aged 11 through 15 years.

2. Rotavirus vaccine (RV).

- The maximum age for the first dose is 14 weeks 6 days. Vaccination should not be initiated for infants aged 15 weeks or older (i.e., 15 weeks 0 days or older).
- Administer the final dose in the series by age 8 months 0 days.
- If Rotarix[®] was administered for the first and second doses, a third dose is not indicated.

3. Diphtheria and tetanus toxoids and acellular pertussis vaccine (DTaP).

- The fifth dose is not necessary if the fourth dose was administered at age 4 years or older.

4. *Haemophilus influenzae* type b conjugate vaccine (Hib).

- Hib vaccine is not generally recommended for persons aged 5 years or older. No efficacy data are available on which to base a recommendation concerning use of Hib vaccine for older children and adults. However, studies suggest good immunogenicity in persons who have sickle cell disease, leukemia, or HIV infection, or who have had a splenectomy; administering 1 dose of Hib vaccine to these persons is not contraindicated.
- If the first 2 doses were PRP-OMP (PedvaxHib[®] or Comvax[®]), and administered at age 11 months or younger, the third (and final) dose should be administered at age 12 through 15 months and at least 8 weeks after the second dose.
- If the first dose was administered at age 7 through 11 months, administer 2 doses separated by 4 weeks and a final dose at age 12 through 15 months.

5. Pneumococcal vaccine.

- Administer 1 dose of pneumococcal conjugate vaccine (PCV) to all healthy children aged 24 through 59 months who have not received at least 1 dose of PCV on or after age 12 months.
- For children aged 24 through 59 months with underlying medical conditions, administer 1 dose of PCV if 3 doses were received previously or administer 2 doses of PCV at least 8 weeks apart if fewer than 3 doses were received previously.
- Administer pneumococcal polysaccharide vaccine (PPSV) to children aged 2 years or older with certain underlying medical conditions (see *MMWR* 2000;49[No. RR-9]), including a cochlear implant, at least 8 weeks after the last dose of PCV.

6. Inactivated poliovirus vaccine (IPV).

- For children who received an all-IPV or all-oral poliovirus (OPV) series, a fourth dose is not necessary if the third dose was administered at age 4 years or older.
- If both OPV and IPV were administered as part of a series, a total of 4 doses should be administered, regardless of the child's current age.

7. Measles, mumps, and rubella vaccine (MMR).

- Administer the second dose at age 4 through 6 years. However, the second dose may be administered before age 4, provided at least 28 days have elapsed since the first dose.
- If not previously vaccinated, administer 2 doses with at least 28 days between doses.

8. Varicella vaccine.

- Administer the second dose at age 4 through 6 years. However, the second dose may be administered before age 4, provided at least 3 months have elapsed since the first dose.
- For persons aged 12 months through 12 years, the minimum interval between doses is 3 months. However, if the second dose was administered at least 28 days after the first dose, it can be accepted as valid.
- For persons aged 13 years and older, the minimum interval between doses is 28 days.

9. Hepatitis A vaccine (HepA).

- HepA is recommended for children older than 1 year who live in areas where vaccination programs target older children or who are at increased risk of infection. See *MMWR* 2006;55(No. RR-7).

10. Tetanus and diphtheria toxoids vaccine (Td) and tetanus and diphtheria toxoids and acellular pertussis vaccine (Tdap).

- Doses of DTaP are counted as part of the Td/Tdap series
- Tdap should be substituted for a single dose of Td in the catch-up series or as a booster for children aged 10 through 18 years; use Td for other doses.

11. Human papillomavirus vaccine (HPV).

- Administer the series to females at age 13 through 18 years if not previously vaccinated.
- Use recommended routine dosing intervals for series catch-up (i.e., the second and third doses should be administered at 2 and 6 months after the first dose). However, the minimum interval between the first and second doses is 4 weeks. The minimum interval between the second and third doses is 12 weeks, and the third dose should be given at least 24 weeks after the first dose.